

Charmful two-body anti-triplet b -baryon decays

Y.K. Hsiao^{1,2,3}, P.Y. Lin⁴, C.C. Lih^{5,2,3}, and C.Q. Geng^{1,2,3}

¹*Chongqing University of Posts & Telecommunications, Chongqing, 400065, China*

²*Physics Division, National Center for Theoretical Sciences, Hsinchu, Taiwan 300*

³*Department of Physics, National Tsing Hua University, Hsinchu, Taiwan 300*

⁴*Department of Electrophysics National Chiao Tung University, Hsinchu, Taiwan 300*

⁵*Department of Optometry, Shu-Zen College of Medicine
and Management, Kaohsiung Hsien, Taiwan 452*

(Dated: November 23, 2015)

Abstract

We study the charmful decays of the two-body $\mathcal{B}_b \rightarrow \mathcal{B}_n M_c$ decays, where $\mathcal{B}_b$ represents the anti-triplet of $(\Lambda_b, \Xi_b^0, \Xi_b^-)$, $\mathcal{B}_n$ stands for the baryon octet and M_c denotes as the charmed meson of $D_{(s)}^{(*)}$, η_c and J/ψ . Explicitly, we predict that $\mathcal{B}(\Lambda_b \rightarrow D_s^- p) = (1.8 \pm 0.3) \times 10^{-5}$, which is within the measured upper bound of $\mathcal{B}(\Lambda_b \rightarrow D_s^- p) < 4.8(5.3) \times 10^{-4}$ at 90% (95%) C.L., and reproduce $\mathcal{B}(\Lambda_b \rightarrow J/\psi \Lambda) = (3.3 \pm 2.0) \times 10^{-4}$ and $\mathcal{B}(\Xi_b^- \rightarrow J/\psi \Xi^-) = (5.1 \pm 3.2) \times 10^{-4}$ in agreement with the data. Moreover, we find that $\mathcal{B}(\Lambda_b \rightarrow \Lambda \eta_c) = (1.5 \pm 0.9) \times 10^{-4}$, $\mathcal{B}(\Xi_b^- \rightarrow \Xi^- \eta_c) = (2.4 \pm 1.5) \times 10^{-4}$ and $\mathcal{B}(\Xi_b^0 \rightarrow \Xi^0 \eta_c, \Xi^0 J/\psi) = (2.3 \pm 1.4, 4.9 \pm 3.0) \times 10^{-4}$, which are accessible to the experiments at the LHCb.

I. INTRODUCTION

The two-body decays of $\Lambda_b \rightarrow \Lambda_c^+ K^-$, $\Lambda_c^+ \pi^-$, $\Lambda_c^+ D^-$, and $\Lambda_c^+ D_s^-$ can be viewed as through the $\Lambda_b \rightarrow \Lambda_c$ transition along with the recoiled mesons K^- , π^- , D_s^- and D^- , respectively, such that one may use the factorization ansatz to get the fractions of the branching ratios as

$$\begin{aligned}\mathcal{R}_{K/\pi} &\equiv \frac{\mathcal{B}(\Lambda_b \rightarrow \Lambda_c^+ K^-)}{\mathcal{B}(\Lambda_b \rightarrow \Lambda_c^+ \pi^-)} \simeq \frac{(|V_{us}|f_K)^2}{(|V_{ud}|f_\pi)^2} = 0.073, \\ \mathcal{R}_{D/D_s} &\equiv \frac{\mathcal{B}(\Lambda_b \rightarrow \Lambda_c^+ D^-)}{\mathcal{B}(\Lambda_b \rightarrow \Lambda_c^+ D_s^-)} \simeq \frac{(|V_{cd}|f_D)^2}{(|V_{cs}|f_{D_s})^2} = 0.034,\end{aligned}\tag{1}$$

which are in agreement with the data, given by [1, 2]

$$\mathcal{R}_{K/\pi} = 0.0731 \pm 0.0016 \pm 0.0016, \quad \mathcal{R}_{D/D_s} = 0.042 \pm 0.003 \pm 0.003.\tag{2}$$

In the same picture, the measured $\mathcal{B}(\Lambda_b \rightarrow pK^-, p\pi)$ can be also explained [3, 4]. In addition, the direct CP violating asymmetry of $\Lambda_b \rightarrow pK^{*-}$ is predicted as large as 20% [5].

On the other hand, the branching ratios of $\Lambda_b \rightarrow D_s^- p$, $\Lambda_b \rightarrow J/\psi \Lambda$ and $\Xi_b^- \rightarrow J/\psi \Xi^-$ are shown as [4, 6]

$$\begin{aligned}\mathcal{B}(\Lambda_b \rightarrow D_s^- p) &= (2.7 \pm 1.4 \pm 0.2 \pm 0.7 \pm 0.1 \pm 0.1) \times 10^{-4} \text{ or} \\ &< 4.8(5.3) \times 10^{-4} \text{ at } 90\% (95\%) \text{ C.L.}, \\ \mathcal{B}(\Lambda_b \rightarrow J/\psi \Lambda) &= (3.0 \pm 1.1) \times 10^{-4}, \\ \mathcal{B}(\Xi_b^- \rightarrow J/\psi \Xi^-) &= (2.0 \pm 0.9) \times 10^{-4},\end{aligned}\tag{3}$$

with $\mathcal{B}(\Lambda_b \rightarrow J/\psi \Lambda)$ and $\mathcal{B}(\Xi_b^- \rightarrow J/\psi \Xi^-)$ converted from the partial observations of $\mathcal{B}(\Lambda_b \rightarrow J/\psi \Lambda)f_{\Lambda_b} = (5.8 \pm 0.8) \times 10^{-5}$ and $\mathcal{B}(\Xi_b^- \rightarrow J/\psi \Xi^-)f_{\Xi_b^-} = (1.02_{-0.21}^{+0.26}) \times 10^{-5}$, where $f_{\Lambda_b} = 0.175 \pm 0.106$ and $f_{\Xi_b} = 0.019 \pm 0.013$ are the fragmentation fractions of the b quark to b -baryons of Λ_b and Ξ_b [7], respectively. Nonetheless, for these $\mathcal{B}_b \rightarrow \mathcal{B}_n M_c$ decays in Eq. (3), the theoretical understanding is still lacking. Since the factorization approach is expected to be reliable in studying the branching ratios of $\mathcal{B}_b \rightarrow \mathcal{B}_n M_c$, in this report, we shall systematically analyze the branching ratios for all possible $\mathcal{B}_b \rightarrow \mathcal{B}_n M_c$ decays, and compare them with the experimental data at the B -factories, as well as the LHCb, where $\mathcal{B}_b$, $\mathcal{B}_n$ and M_c correspond to the anti-triplet b -baryon of $(\Lambda_b, \Xi_b^0, \Xi_b^-)$, baryon octet and charmed meson, respectively.




FIG. 1. Diagrams for two-body charmful $\mathcal{B}_b \rightarrow \mathcal{B}_n M_c$ decays.

II. FORMALISM

As the studies in Refs. [8–14], based on the factorization approach, the amplitudes for the two-body charmful b -baryon decays are presented in terms of the decaying process of the $\mathcal{B}_b \rightarrow \mathcal{B}_n$ transition with the recoiled charmed meson M_c . From Fig. 1a, the amplitudes of $\mathcal{B}_b \rightarrow \mathcal{B}_n M_c$ via the quark-level $b \rightarrow u \bar{c} q$ transition are factorized as

$$\mathcal{A}_1(\mathcal{B}_b \rightarrow \mathcal{B}_n M_c) = \frac{G_F}{\sqrt{2}} V_{ub} V_{cq}^* a_1 \langle M_c | \bar{q} \gamma^\mu (1 - \gamma_5) c | 0 \rangle \langle \mathcal{B}_n | \bar{u} \gamma_\mu (1 - \gamma_5) b | \mathcal{B}_b \rangle, \quad (4)$$

where G_F is the Fermi constant, $V_{ub, cq}$ are the CKM matrix elements, while the explicit decay modes are

$$\Lambda_b \rightarrow p M_c, \quad \Xi_b^- \rightarrow \Lambda(\Sigma^0) M_c, \quad \Xi_b^0 \rightarrow \Sigma^+ M_c \quad (5)$$

with $q = d(s)$ for $M_c = D^{(*)-}(D_s^{(*)-})$. On the other hand, the amplitudes via the quark-level $b \rightarrow c \bar{u} q$ ($b \rightarrow c \bar{c} q$) transition in Fig. 1b can be written as

$$\mathcal{A}_2(\mathcal{B}_b \rightarrow \mathcal{B}_n M_c) = \frac{G_F}{\sqrt{2}} V_{cb} V_{q_1 q}^* a_2 \langle M_c | \bar{c} \gamma^\mu (1 - \gamma_5) q_1 | 0 \rangle \langle \mathcal{B}_n | \bar{q} \gamma_\mu (1 - \gamma_5) b | \mathcal{B}_b \rangle, \quad (6)$$

with $q_1 = u$ for $M_c = D^{(*)0}$ and $q_1 = c$ for $M_c = \eta_c$ and J/ψ , where the decays of $\mathcal{B}_b \rightarrow \mathcal{B}_n M_c$ are

$$\begin{aligned} \Lambda_b \rightarrow n M_c, \quad \Xi_b^- \rightarrow \Sigma^- M_c, \quad \Xi_b^0 \rightarrow \Lambda(\Sigma^0) M_c \quad \text{for } q_2 = d, \\ \Lambda_b \rightarrow \Lambda(\Sigma^0) M_c, \quad \Xi_b^- \rightarrow \Xi^- M_c, \quad \Xi_b^0 \rightarrow \Xi^0 M_c \quad \text{for } q_2 = s. \end{aligned} \quad (7)$$

In this study, we will exclude the study of $\Lambda_b \rightarrow n M_c$ due to the elusive neutron in the B -factories. The amplitudes $\mathcal{A}_{1,2}$ via the W -boson exchange diagrams are led to be the color-allowed and color-suppressed processes. The parameters a_1 and a_2 in Eqs. (4) and (6)

are presented as [15, 16]

$$a_1 = c_1^{eff} + \frac{c_2^{eff}}{N_c}, a_2 = c_2^{eff} + \frac{c_1^{eff}}{N_c}, \quad (8)$$

with the effective Wilson coefficients $(c_1^{eff}, c_2^{eff}) = (1.168, -0.365)$, respectively, where the color number N_c should be taken as a floating number from $2 \rightarrow \infty$ to account for the non-factorizable effects in the generalized factorization instead of $N_c = 3$. The matrix elements for $P_c = (\eta_c, D)$ and $V_c = J(/\psi, D^*)$ productions read

$$\begin{aligned} \langle P_c | A_\mu^c | 0 \rangle &= -i f_{P_c} q_\mu, \\ \langle V_c | V_\mu^c | 0 \rangle &= m_{V_c} f_{V_c} \varepsilon_\mu^*, \end{aligned} \quad (9)$$

with $V_\mu^c(A_\mu^c) = \bar{q}\gamma^\mu(\gamma_5)c$ or $\bar{c}\gamma^\mu(\gamma_5)q_1$, where q_μ and ε_μ^* are the four-momentum and polarization, respectively. Those of the $\mathcal{B}_b \rightarrow \mathcal{B}_n$ baryon transition in Eq. (4) have the general forms:

$$\begin{aligned} \langle \mathcal{B}_n | \bar{q}\gamma_\mu b | \mathcal{B}_b \rangle &= \bar{u}_{\mathcal{B}_n} \left[f_1 \gamma_\mu + \frac{f_2}{m_{\mathcal{B}_b}} i \sigma_{\mu\nu} q^\nu + \frac{f_3}{m_{\mathcal{B}_b}} q_\mu \right] u_{\mathcal{B}_b}, \\ \langle \mathcal{B}_n | \bar{q}\gamma_\mu \gamma_5 b | \mathcal{B}_b \rangle &= \bar{u}_{\mathcal{B}_n} \left[g_1 \gamma_\mu + \frac{g_2}{m_{\mathcal{B}_b}} i \sigma_{\mu\nu} q^\nu + \frac{g_3}{m_{\mathcal{B}_b}} q_\mu \right] \gamma_5 u_{\mathcal{B}_b}, \end{aligned} \quad (10)$$

where f_j (g_j) ($j = 1, 2, 3$) are the form factors. We are able to relate the different $\mathcal{B}_b \rightarrow \mathcal{B}_n$ transition form factors based on the $SU(3)$ flavor and $SU(2)$ spin symmetries, which have been used to connect the space-like $\mathcal{B}_n \rightarrow \mathcal{B}'_n$ transition form factors in the neutron decays [17], and the time-like $0 \rightarrow \mathcal{B}_n \bar{\mathcal{B}}'_n$ baryonic form factors as well as the $B \rightarrow \mathcal{B}_n \bar{\mathcal{B}}'_n$ transition form factors in the baryonic B decays [18–22]. Specifically, $V_\mu^q = \bar{q}\gamma_\mu b$ and $A_\mu^q = \bar{q}\gamma_\mu \gamma_5 b$ as the two currents in Eq. (10) can be combined as the right-handed chiral current, that is, $J_{\mu,R}^q = (V_\mu^q + A_\mu^q)/2$. Consequently, we have [17]:

$$\langle \mathcal{B}_{n,\uparrow+\downarrow} | J_{\mu,R}^q | \mathcal{B}_{b,\uparrow+\downarrow} \rangle = \bar{u}_{\mathcal{B}_n} \left[\gamma_\mu \frac{1+\gamma_5}{2} G^\uparrow(q^2) + \gamma_\mu \frac{1-\gamma_5}{2} G^\downarrow(q^2) \right] u_{\mathcal{B}_b}, \quad (11)$$

where the baryon helicity states $|\mathcal{B}_{n(b),\uparrow+\downarrow}\rangle \equiv |\mathcal{B}_{n(b),\uparrow}\rangle + |\mathcal{B}_{n(b),\downarrow}\rangle$ are regarded as the baryon chiral states $|\mathcal{B}_{n(b),R+L}\rangle$ at the large momentum transfer, while $G^\uparrow(q^2)$ and $G^\downarrow(q^2)$ are the right-handed and left-handed form factors, defined by

$$G^\uparrow(q^2) = e_\parallel^\uparrow G_\parallel(q^2) + e_\parallel^\uparrow G_\parallel(q^2), \quad G^\downarrow(q^2) = e_\parallel^\downarrow G_\parallel(q^2) + e_\parallel^\downarrow G_\parallel(q^2), \quad (12)$$

with the constants $e_{||(\overline{||})}^\uparrow$ and $e_{||(\overline{||})}^\downarrow$ to sum over the chiral charges via the $\mathcal{B}_b \rightarrow \mathcal{B}_n$ transition, given by

$$\begin{aligned} e_{||}^\uparrow &= \langle \mathcal{B}_{n,\uparrow} | \mathbf{Q}_{||} | \mathcal{B}_{b,\uparrow} \rangle, & e_{||}^\downarrow &= \langle \mathcal{B}_{n,\uparrow} | \mathbf{Q}_{||} | \mathcal{B}_{b,\downarrow} \rangle, \\ e_{\overline{||}}^\uparrow &= \langle \mathcal{B}_{n,\uparrow} | \mathbf{Q}_{\overline{||}} | \mathcal{B}_{b,\uparrow} \rangle, & e_{\overline{||}}^\downarrow &= \langle \mathcal{B}_{n,\uparrow} | \mathbf{Q}_{\overline{||}} | \mathcal{B}_{b,\downarrow} \rangle, \\ e_{||}^\downarrow &= \langle \mathcal{B}_{n,\downarrow} | \mathbf{Q}_{||} | \mathcal{B}_{b,\downarrow} \rangle, & e_{\overline{||}}^\uparrow &= \langle \mathcal{B}_{n,\downarrow} | \mathbf{Q}_{\overline{||}} | \mathcal{B}_{b,\downarrow} \rangle. \end{aligned} \quad (13)$$

Note that $\mathbf{Q}_{||(\overline{||})} = \sum_i Q_{||(\overline{||})}(i)$ with $i = 1, 2, 3$ as the the chiral charge operators are from $Q_R^q \equiv J_{0,R}^q = q_R^\dagger b_R$, converting the b quark in $|\mathcal{B}_{b,\uparrow,\downarrow}\rangle$ into the q one, while the converted q quark can be parallel or antiparallel to the $\mathcal{B}_b$'s helicity, denoted as the subscript ($||$ or $\overline{||}$). By comparing Eq. (10) with Eqs. (11), (12), and (13), we obtain

$$\begin{aligned} f_1 &= (e_{||}^\uparrow + e_{||}^\downarrow)G_{||} + (e_{\overline{||}}^\uparrow + e_{\overline{||}}^\downarrow)G_{\overline{||}}, \\ g_1 &= (e_{||}^\uparrow - e_{||}^\downarrow)G_{||} + (e_{\overline{||}}^\uparrow - e_{\overline{||}}^\downarrow)G_{\overline{||}}, \end{aligned} \quad (14)$$

with $f_{2,3} = 0$ and $g_{2,3} = 0$ due to the helicity conservation, as those derived in Refs. [8, 10, 23]. It is interesting to see that, as the helicity-flip terms, the theoretical calculations from the loop contributions to $f_{2,3}$ ($g_{2,3}$) indeed result in the values to be one order of magnitude smaller than that of $f_1(g_1)$, which can be safely neglected. In the double-pole momentum dependences, f_1 and g_1 can be given as [3]

$$f_1(q^2) = \frac{f_1(0)}{(1 - q^2/m_{\mathcal{B}_b}^2)^2}, \quad g_1(q^2) = \frac{g_1(0)}{(1 - q^2/m_{\mathcal{B}_b}^2)^2}, \quad (15)$$

such that it is reasonable to parameterize the chiral form factors to be $(1 - q^2/m_{\mathcal{B}_b}^2)^2 G_{||(\overline{||})} = C_{||(\overline{||})}$. Subsequently, from

$$\begin{aligned} (e_{||}^\uparrow, e_{||}^\downarrow, e_{\overline{||}}^\uparrow, e_{\overline{||}}^\downarrow) &= (-\sqrt{3/2}, 0, 0, 0) \text{ for } \langle p | J_{\mu,R}^u | \Lambda_b \rangle, \\ (e_{||}^\uparrow, e_{||}^\downarrow, e_{\overline{||}}^\uparrow, e_{\overline{||}}^\downarrow) &= (1, 0, 0, 0) \text{ for } \langle \Lambda | J_{\mu,R}^s | \Lambda_b \rangle, \\ (e_{||}^\uparrow, e_{||}^\downarrow, e_{\overline{||}}^\uparrow, e_{\overline{||}}^\downarrow) &= (0, 0, 0, 0) \text{ for } \langle \Sigma^0 | J_{\mu,R}^s | \Lambda_b \rangle, \end{aligned} \quad (16)$$

we get $f_1(0) = g_1(0) = -\sqrt{3/2} C_{||}$ for $\langle p | \bar{u} \gamma_\mu (\gamma_5) b | \Lambda_b \rangle$, $f_1(0) = g_1(0) = C_{||}$ for $\langle \Lambda | \bar{s} \gamma_\mu (\gamma_5) b | \Lambda_b \rangle$, and $f_1(0) = g_1(0) = 0$ for $\langle \Sigma^0 | \bar{s} \gamma_\mu (\gamma_5) b | \Lambda_b \rangle$, similar to the results based on the heavy-quark and large-energy symmetries in Ref. [23] for the $\Lambda_b \rightarrow (p, \Lambda, \Sigma)$ transitions. When we further extend the study to the anti-triplet b -baryons: $(\Xi_b^-, \Xi_b^0, \Lambda_b^0)$ shown in Table I, we find that the relation of $f_1 = g_1$ is uniquely determined for the anti-triplet b -baryon transitions.

TABLE I. Relations between the transition matrix elements.

$\langle \mathcal{B}_n (\bar{q}b) \mathcal{B}_b \rangle$	$f_1(0) = g_1(0)$
$\langle p (\bar{u}b) \Lambda_b \rangle$	$-\sqrt{\frac{3}{2}}C_{ }$
$\langle \Lambda (\bar{u}b) \Xi_b^- \rangle$	$\frac{1}{2}C_{ }$
$\langle \Sigma^0 (\bar{u}b) \Xi_b^- \rangle$	$-\sqrt{\frac{3}{4}}C_{ }$
$\langle \Sigma^+ (\bar{u}b) \Xi_b^0 \rangle$	$-\sqrt{\frac{3}{2}}C_{ }$
$\langle \Sigma^- (\bar{d}b) \Xi_b^- \rangle$	$\sqrt{\frac{3}{2}}C_{ }$
$\langle \Lambda (\bar{d}b) \Xi_b^0 \rangle$	$-\frac{1}{2}C_{ }$
$\langle \Sigma^0 (\bar{d}b) \Xi_b^0 \rangle$	$\sqrt{\frac{3}{4}}C_{ }$
$\langle \Lambda (\bar{s}b) \Lambda_b \rangle$	$C_{ }$
$\langle \Sigma^0 (\bar{s}b) \Lambda_b \rangle$	0
$\langle \Xi^- (\bar{s}b) \Xi_b^- \rangle$	$\sqrt{\frac{3}{2}}C_{ }$
$\langle \Xi^0 (\bar{s}b) \Xi_b^0 \rangle$	$-\sqrt{\frac{3}{2}}C_{ }$

III. NUMERICAL RESULTS AND DISCUSSIONS

For the numerical analysis, the CKM matrix elements in the Wolfenstein parameterization taken from the PDG [4] are given by

$$\begin{aligned}
 (V_{ub}, V_{cb}) &= (A\lambda^3(\rho - i\eta), A\lambda^2), \\
 (V_{cd} = -V_{us}, V_{cs} = V_{ud}) &= (-\lambda, 1 - \lambda^2/2),
 \end{aligned} \tag{17}$$

with $(\lambda, A, \rho, \eta) = (0.225, 0.814, 0.120 \pm 0.022, 0.362 \pm 0.013)$. The meson decay constants are adopted as $(f_{\eta_c}, f_{J/\psi}) = (387 \pm 7, 418 \pm 9)$ MeV [24], $(f_D, f_{D_s}) = (204.6 \pm 5.0, 257.5 \pm 4.6)$ MeV [4], and $(f_{D^*}, f_{D_s^*}) = (252.2 \pm 22.7, 305.5 \pm 27.3)$ MeV [25]. As given in Ref. [3] to explain the branching ratios and CP violating asymmetries of $\Lambda_b \rightarrow p(K^-, \pi^-)$, we have $|\sqrt{3/2}C_{||}| = 0.136 \pm 0.009$ for $\langle p | \bar{u}\gamma_\mu(\gamma_5)b | \Lambda_b \rangle$, which is consistent with the value of 0.14 ± 0.03 in the light-cone sum rules [23] and the theoretical calculations in Refs. [8, 10]. With $\mathcal{B}(\Lambda_b \rightarrow J/\psi\Lambda)$ and $\mathcal{B}(\Xi_b^- \rightarrow J/\psi\Xi^-)$ in Eq. (3) as the experimental inputs, we can estimate the non-factorizable effects by deviating the color number $N_c = 3$ to be between 2 to ∞ , such that we obtain $N_c = 2.15 \pm 0.17$, representing controllable non-factorizable effects [26] with $(a_1, a_2) = (1.00 \pm 0.01, 0.18 \pm 0.05)$ from Eq. (8). We list the branching

ratios of all possible two-body anti-triplet b -baryon decays in Table II and Table III, where the uncertainties are fitted with those from (ρ, η, N_c) , the decay constants and $|\sqrt{3/2}C_{||}|$.

The decay branching ratios in Table II are given by a_1 with $N_c = (2.15 \pm 0.17, \infty)$ as the theoretical inputs to demonstrate the insensitive non-factorizable effects. Note that $N_c = 2.15 \pm 0.17$ is fitted from $\mathcal{B}(\Lambda_b \rightarrow J/\psi \Lambda)$ and $\mathcal{B}(\Xi_b^- \rightarrow J/\psi \Xi^-)$, while $N_c = \infty$ results in $a_1 \simeq c_1^{eff}$, wildly used in the generalized factorization. As the first measurement for the color-allowed decay mode, the predicted $\mathcal{B}(\Lambda_b \rightarrow D_s^- p) = (1.8 \pm 0.3) \times 10^{-5}$ or $(2.5 \pm 0.4) \times 10^{-5}$ in Table II seems to disagree with the data in Eq. (3). Nonetheless, the predicted numbers driven by a_1 can be reliable as it is insensitive to the non-factorizable effects, whereas the data with the upper bound has a large uncertainty. Despite the color-allowed modes, the decay branching ratios of $D^{(*)-}$ are found to be 30 times smaller than the $D_s^{(*)-}$ counterparts. This can be simply understood by the relation of $(V_{cd}/V_{cs})^2(f_{D^{(*)}}/f_{D_s^{(*)}})^2 \simeq 0.03$. It is also interesting to note that the vector meson modes are 2 times as large as their pseudoscalar meson counterparts.

For the decay modes driven by a_2 as shown in Table III, we only list the results with $a_2 = 0.18 \pm 0.05$ ($N_c = 2.15 \pm 0.17$). The reason is that $a_2 \simeq c_2^{eff} = -0.365$ with $N_c = \infty$ yields $\mathcal{B}(\Lambda_b \rightarrow J/\psi \Lambda) = (1.4 \pm 0.2) \times 10^{-3}$ and $\mathcal{B}(\Xi_b^- \rightarrow J/\psi \Xi^-) = (2.1 \pm 0.3) \times 10^{-3}$, which are in disagreement with the data in Eq. (3), demonstrating that the decays are sensitive

TABLE II. The branching ratios of all possible two-body anti-triplet b -baryon decays with a_1 fitted by $N_c = (2.15 \pm 0.17, \infty)$.

$M_c =$	D^-	D^{*-}
$\mathcal{B}(\Lambda_b \rightarrow p M_c)$	$(6.0 \pm 1.0, 8.2 \pm 1.4) \times 10^{-7}$	$(1.2 \pm 0.3, 1.6 \pm 0.4) \times 10^{-6}$
$\mathcal{B}(\Xi_b^- \rightarrow \Lambda M_c)$	$(1.1 \pm 0.2, 1.5 \pm 0.2) \times 10^{-7}$	$(2.2 \pm 0.6, 3.0 \pm 0.8) \times 10^{-7}$
$\mathcal{B}(\Xi_b^- \rightarrow \Sigma^0 M_c)$	$(3.3 \pm 0.5, 4.5 \pm 0.7) \times 10^{-7}$	$(6.6 \pm 1.6, 9.0 \pm 2.2) \times 10^{-7}$
$\mathcal{B}(\Xi_b^0 \rightarrow \Sigma^+ M_c)$	$(6.3 \pm 1.0, 8.6 \pm 1.4) \times 10^{-7}$	$(1.3 \pm 0.3, 1.7 \pm 0.4) \times 10^{-6}$
$M_c =$	D_s^-	D_s^{*-}
$\mathcal{B}(\Lambda_b \rightarrow p M_c)$	$(1.8 \pm 0.3, 2.5 \pm 0.4) \times 10^{-5}$	$(3.5 \pm 0.9, 4.7 \pm 1.2) \times 10^{-5}$
$\mathcal{B}(\Xi_b^- \rightarrow \Lambda M_c)$	$(3.4 \pm 0.5, 4.6 \pm 0.7) \times 10^{-6}$	$(6.4 \pm 1.6, 8.8 \pm 2.2) \times 10^{-6}$
$\mathcal{B}(\Xi_b^- \rightarrow \Sigma^0 M_c)$	$(9.9 \pm 1.5, 13.6 \pm 2.1) \times 10^{-6}$	$(1.9 \pm 0.5, 2.6 \pm 0.6) \times 10^{-5}$
$\mathcal{B}(\Xi_b^0 \rightarrow \Sigma^+ M_c)$	$(1.9 \pm 0.3, 2.6 \pm 0.4) \times 10^{-5}$	$(3.6 \pm 0.9, 4.9 \pm 1.2) \times 10^{-5}$

to the non-factorizable effects. From Table III, we see that both $\mathcal{B}(\Lambda_b \rightarrow J/\psi \Lambda)$ and $\mathcal{B}(\Xi_b^- \rightarrow J/\psi \Xi^-)$ are reproduced to agree with the data in Eq. (3) within errors. Note that $\mathcal{B}(\Lambda_b \rightarrow J/\psi \Lambda)/\mathcal{B}(\Xi_b^- \rightarrow J/\psi \Xi^-) \simeq 0.65$ in our calculation results from $(C_{||})^2/(\sqrt{3/2}C_{||})^2 \simeq 0.67$ as the ratio of their form factors in Table I, which is in accordance with the $SU(3)$ flavor and $SU(2)$ spin symmetries. The more precise measurement of this ratio in the future will test the validity of the symmetries. As $\mathcal{B}(\Xi_b^- \rightarrow J/\psi \Xi^-) = O(10^{-4})$, we emphasize that more experimental searches should be done for the two-body Ξ_b decays, while most of the recent observations are from the Λ_b decays. Since $\mathcal{B}(\Lambda_b \rightarrow \Sigma^0 M_c) = 0$ results from the $SU(3)$ flavor and $SU(2)$ spin symmetries as well as the heavy flavor symmetry, its non-zero measurement will imply for the broken symmetry. Through the $b \rightarrow c\bar{c}s$ transition at the quark level, $\mathcal{B}(\Lambda_b \rightarrow \Lambda M_c)$, $\mathcal{B}(\Xi_b^- \rightarrow \Xi^- M_c)$ and $\mathcal{B}(\Xi_b^0 \rightarrow \Xi^0 M_c)$ with $M_c = \eta_c$ and J/ψ are all $O(10^{-4})$ as shown in the right-bottom of Table II. In contrast, the neutral $D^{(*)0}$ modes

TABLE III. The branching ratios of all possible two-body anti-triplet b -baryon decays with a_2 fitted by $N_c = 2.15 \pm 0.17$.

$M_c =$	D^0	D^{*0}
$\mathcal{B}(\Xi_b^- \rightarrow \Sigma^- M_c)$	$(5.3 \pm 3.3) \times 10^{-5}$	$(1.1 \pm 0.7) \times 10^{-4}$
$\mathcal{B}(\Xi_b^0 \rightarrow \Lambda^0 M_c)$	$(8.6 \pm 5.3) \times 10^{-6}$	$(1.7 \pm 1.1) \times 10^{-5}$
$\mathcal{B}(\Xi_b^0 \rightarrow \Sigma^0 M_c)$	$(2.5 \pm 1.6) \times 10^{-5}$	$(5.0 \pm 3.4) \times 10^{-5}$
$\mathcal{B}(\Lambda_b \rightarrow \Lambda M_c)$	$(1.6 \pm 1.0) \times 10^{-6}$	$(3.3 \pm 2.2) \times 10^{-6}$
$\mathcal{B}(\Lambda_b \rightarrow \Sigma^0 M_c)$	0	0
$\mathcal{B}(\Xi_b^- \rightarrow \Xi^- M_c)$	$(2.7 \pm 1.7) \times 10^{-6}$	$(5.5 \pm 3.6) \times 10^{-6}$
$\mathcal{B}(\Xi_b^0 \rightarrow \Xi^0 M_c)$	$(2.6 \pm 1.6) \times 10^{-6}$	$(5.2 \pm 3.5) \times 10^{-6}$
$M_c =$	η_c	J/ψ
$\mathcal{B}(\Xi_b^- \rightarrow \Sigma^- M_c)$	$(1.4 \pm 0.8) \times 10^{-5}$	$(2.9 \pm 1.8) \times 10^{-5}$
$\mathcal{B}(\Xi_b^0 \rightarrow \Lambda^0 M_c)$	$(2.3 \pm 1.4) \times 10^{-6}$	$(4.7 \pm 2.9) \times 10^{-6}$
$\mathcal{B}(\Xi_b^0 \rightarrow \Sigma^0 M_c)$	$(6.6 \pm 4.1) \times 10^{-6}$	$(1.4 \pm 0.8) \times 10^{-5}$
$\mathcal{B}(\Lambda_b \rightarrow \Lambda M_c)$	$(1.5 \pm 0.9) \times 10^{-4}$	$(3.3 \pm 2.0) \times 10^{-4}$
$\mathcal{B}(\Lambda_b \rightarrow \Sigma^0 M_c)$	0	0
$\mathcal{B}(\Xi_b^- \rightarrow \Xi^- M_c)$	$(2.4 \pm 1.5) \times 10^{-4}$	$(5.1 \pm 3.2) \times 10^{-4}$
$\mathcal{B}(\Xi_b^0 \rightarrow \Xi^0 M_c)$	$(2.3 \pm 1.4) \times 10^{-4}$	$(4.9 \pm 3.0) \times 10^{-4}$

via the $b \rightarrow c\bar{c}d$ transition have the branching ratios of order 10^{-6} caused by the suppression of $(V_{cb}V_{cd})^2/(V_{cb}V_{cs})^2 \simeq 0.225^2$. Finally, we remark that $\mathcal{B}(\Xi_b^- \rightarrow \Xi^- M_c) \simeq \mathcal{B}(\Xi_b^0 \rightarrow \Xi^0 M_c)$ is due to the isospin symmetry.

IV. CONCLUSIONS

In sum, we have studied all possible anti-triplet $\mathcal{B}_b$ decays of the two-body charmed $\mathcal{B}_b \rightarrow \mathcal{B}_n M_c$ decays. We have found $\mathcal{B}(\Lambda_b \rightarrow D_s^- p) = (1.8 \pm 0.3) \times 10^{-5}$, which is within the measured upper bound of $\mathcal{B}(\Lambda_b \rightarrow D_s^- p) < 4.8(5.3) \times 10^{-4}$ at 90% (95%) C.L., and reproduced $\mathcal{B}(\Lambda_b \rightarrow J/\psi \Lambda) = (3.3 \pm 2.0) \times 10^{-4}$ and $\mathcal{B}(\Xi_b^- \rightarrow J/\psi \Xi^-) = (5.1 \pm 3.2) \times 10^{-4}$ in agreement with the data. Moreover, we have predicted $\mathcal{B}(\Lambda_b \rightarrow \Lambda \eta_c) = (1.5 \pm 0.9) \times 10^{-4}$, $\mathcal{B}(\Xi_b^- \rightarrow \Xi^- \eta_c) = (2.4 \pm 1.5) \times 10^{-4}$, and $\mathcal{B}(\Xi_b^0 \rightarrow \Xi^0 \eta_c, \Xi^0 J/\psi) = (2.3 \pm 1.4, 4.9 \pm 3.0) \times 10^{-4}$, which are accessible to the experiments at the LHCb, while $\mathcal{B}(\Xi_b^- \rightarrow \Xi^- M_c) \simeq \mathcal{B}(\Xi_b^0 \rightarrow \Xi^0 M_c)$ is due to the isospin symmetry.

ACKNOWLEDGMENTS

The work was supported in part by National Center for Theoretical Science, National Science Council (NSC-101-2112-M-007-006-MY3 and NSC-102-2112-M-471-001-MY3), MoST (MoST-104-2112-M-007-003-MY3) and National Tsing Hua University (104N2724E1).

-
- [1] R. Aaij *et al.* [LHCb Collaboration], Phys. Rev. D **89**, 032001 (2014).
 - [2] R. Aaij *et al.* [LHCb Collaboration], Phys. Rev. Lett. **112**, 202001 (2014).
 - [3] Y.K. Hsiao and C.Q. Geng, Phys. Rev. D **91**, 116007 (2015).
 - [4] K.A. Olive *et al.* [Particle Data Group Collaboration], Chin. Phys. C **38**, 090001 (2014).
 - [5] C.Q. Geng, Y.K. Hsiao and J.N. Ng, Phys. Rev. Lett. **98**, 011801 (2007).
 - [6] R. Aaij *et al.* [LHCb Collaboration], JHEP **1404**, 087 (2014).
 - [7] Y.K. Hsiao, P.Y. Lin, L.W. Luo, and C.Q. Geng, to be published in Phys. Lett. B (2015), arXiv:1510.01808 [hep-ph].
 - [8] T. Gutsche *et. al.*, Phys. Rev. D **88**, 114018 (2013).
 - [9] Y. Liu, X.H. Guo and C. Wang, Phys. Rev. D **91**, 016006 (2015).

- [10] Z.T. Wei, H.W. Ke and X.Q. Li, Phys. Rev. D **80**, 094016 (2009).
- [11] Fayyazuddin and Riazuddin, Phys. Rev. D **58**, 014016 (1998).
- [12] C.H. Chou, H.H. Shih, S.C. Lee and H.n. Li, Phys. Rev. D **65**, 074030 (2002).
- [13] M.A. Ivanov *et. al.*, Phys. Rev. D **57**, 5632 (1998).
- [14] H.Y. Cheng, Phys. Rev. D **56**, 2799 (1997).
- [15] Y.H. Chen *et al.*, Phys. Rev. **D60**, 094014 (1999); H.Y. Cheng and K.C. Yang, *ibid*, **D62**, 054029 (2000).
- [16] A. Ali, G. Kramer, and C.D. Lu, Phys. Rev. **D58**, 094009 (1998).
- [17] G.P. Lepage and S.J. Brodsky, Phys. Rev. Lett. **43**, 545(1979) [Erratum-*ibid.* **43**, 1625 (1979)].
- [18] C.K. Chua, W.S. Hou and S.Y. Tsai, Phys. Rev. D **66**, 054004 (2002).
- [19] C.K. Chua and W.S. Hou, Eur. Phys. J. C **29**, 27 (2003).
- [20] C.H. Chen, H.Y. Cheng, C.Q. Geng and Y.K. Hsiao, Phys. Rev. D **78**, 054016 (2008).
- [21] C.Q. Geng and Y.K. Hsiao, Phys. Rev. D **85**, 017501 (2012).
- [22] Y.K. Hsiao and C.Q. Geng, Phys. Rev. D **91**, 077501 (2015).
- [23] A. Khodjamirian *et al.*, JHEP **1109**, 106 (2011); T. Mannel and Y. M. Wang, JHEP **1112**, 067 (2011).
- [24] D. Becirevic *et. al.*, Nucl. Phys. B **883**, 306 (2014).
- [25] W. Lucha, D. Melikhov and S. Simula, Phys. Lett. B **735**, 12 (2014).
- [26] M. Neubert and A.A. Petrov, Phys. Rev. D **68**, 014004 (2003).